

CASSINI TOST SEGMENT

Rev 159 Handoff Package

Segment Boundary 2012-002T00:22:00 - 2012-003T18:23:00

25 Jan 2011 (SMT updated 16 May 2011)

Deb Chattopadhyay

SMT report and SPASS

Science Highlights

Notes & Liens

Integration Checklist

SMT report

Data volume summary Updated 5/16/11

DATA VOLUME SUMMARY --- TRANSFER FRAME OVERHEAD INCLUDED (80 BITS PER 8800-BIT FRAME)

			OBSERVATION_PERIOD					DOWNLINK_PASS							 		
			 P4 P5 				 P5 	 RECC 	RDED	PLAYBACK							
DOWNLINK PASS NAME	Start doy hh:mm	End doy hh:mm	START	SCI (Mb)	HK+E (Mb)	TOTAL	CPACTY (Mb)	MRGN (Mb)	 OPNAV (Mb)	 SCI (Mb)	ENGR (Mb)	 TOTAL (Mb)	CPACTY (Mb)	MARGN (Mb)	NET_MA	ARGN (%)	CAROVR
SP_159EA_G70METNON003_PRIME	003 08:23	003 18:23	0	2661 	136 	2797	3322 	526	0	316	59	3171 	3171	-1	0	0% 	 0

SPASS

Request	Riders	Start (SCET)	Start (Epoch)	Duration	End (SCET)	Primary	Secondary	Comments
Sequence S71, length = 70 days		2011-320T03:02:00		069T19:53:00	2012-024T22:55:00			
Titan Flyby T80 Segment		2012-002T00:22:00		001T18:01:00	2012-003T18:23:00			
SP_159TI_WAYPTTURN002_PRIME		2012-002T00:22:00		000T00:40:00	2012-002T01:02:00	NEG_Y to Titan	POS_X to 231.1/-23.9	
NEW WAYPOINT		2012-002T01:02:00		001T05:51:00	2012-003T06:53:00	NEG_Y to Titan	POS_X to 231.1/-23.9	
SP_159TI_DEADTIME002_PRIME		2012-002T01:02:00		000T00:15:00	2012-002T01:17:00	NEG_Y to Titan	POS_X to 231.1/-23.9	
CIRS_159TI_FIRNADCMP001_PRIME	I, V	2012-002T01:17:00	GMB_E159_TITAN_T80-000T13:56:38	000T04:56:38	2012-002T06:13:38	CIRS_FP1 to Titan	PIC	
CIRS_159TI_MIRLMBMAP001_PRIME	I, V	2012-002T06:13:38	GMB_E159_TITAN_T80-000T09:00:00	000T04:00:00	2012-002T10:13:38	CIRS_FPB to Titan	PIC	
CIRS_159TI_FIRNADMAP001_PRIME	I, V	2012-002T10:13:38	GMB_E159_TITAN_T80-000T05:00:00	000T02:45:00	2012-002T12:58:38	CIRS_FP1 to Titan	PIC	
ISS_159TI_HIGHRES001_PRIME	C, M, V	2012-002T12:58:38	GMB_E159_TITAN_T80-000T02:15:00	000T04:30:00	2012-002T17:28:38	ISS_NAC to Titan	POS_X to 231.1/-23.9	
159TI (t) T80 TITAN Inboun		2012-002T15:13:38		000T00:00:01	2012-002T15:13:39			
RADAR_159TI_T80OUTRAD001_PRIME		2012-002T17:28:38	GMB_E159_TITAN_T80+000T02:15:00	000T02:45:00	2012-002T20:13:38	NEG_Z to Titan	NEG_X to NTP	Use -X to NTP and -Y to NTP for
								the two polarizations.
VIMS_159TI_STELLOCC001_PRIME	С		GMB_E159_TITAN_T80+000T05:00:00			VIMS_IR to Titan	NEG_X to Sun	
VIMS_159TI_GLOBMAP001_PRIME	C, I	2012-003T00:13:38	GMB_E159_TITAN_T80+000T09:00:00	000T05:44:22	2012-003T05:58:00	VIMS_IR to Titan	NEG_X to Sun	
SP_159NA_DEADTIME003_PRIME		2012-003T05:58:00	GMB_E159_TITAN_T80+000T14:44:22	000T00:15:00	2012-003T06:13:00	NEG_Y to Titan	POS_X to 231.1/-23.9	
SP_159EA_DLTURN003_PRIME		2012-003T06:13:00		000T00:40:00	2012-003T06:53:00	XBAND to Earth	NEG_X to NEP	
NEW WAYPOINT		2012-003T06:53:00		000T11:30:00	2012-003T18:23:00	XBAND to Earth	NEG_X to NEP	
SP_159EA_YBIAS003_PRIME	E	2012-003T06:53:00		000T01:30:00	2012-003T08:23:00	XBAND to Earth	NEG_X to NEP	
SP_159EA_G70METNON003_PRIME	C	2012-003T08:23:00		000T10:00:00	2012-003T18:23:00	XBAND to Earth	NEG_X to NEP	NEG_X to NEP or NSP, CAPS

Science Highlights

Jan 2 (DOY 002) - As the lead instrument on the inbound approach to Titan, CIRS views Titan at various distances and spectral resolutions to make maps of atmospheric and surface temperature. On this high-altitude encounter, ISS will perform high-resolution observations around closest-approach along the anti-Saturnian and trailing hemispheres at mid- to high southern latitudes. After c/a RADAR will do radiometry of the subsaturn hemisphere. Outbound, VIMS will perform stellar occultations of CW Leo and R Leo which will constrain the composition and the spectral properties of Titan's atmosphere.

During the flyby, RPWS will measure thermal plasmas in Titan's ionosphere and surrounding environment; search for lightning in Titan's atmosphere; investigate the interaction of Titan with Saturn's magnetosphere. CAPS will Measure the ion and electron temperatures, densities and the ion composition and flow field in the vicinity of Titan, to characterize and understand its interaction with the magnetosphere of Saturn. Observe any seasonal or other long-term variability in the characteristics of this interaction, and study the processes by which Titan's atmosphere and ionosphere are lost to the magnetosphere.

Jan 3 (DOY 003) – VIMS will perform a global mapping observation of the high southern latitudes. ISS will ride along with VIMS' outbound observations of Titan's leading hemisphere, including Senkyo. This is one of the last few views of the high southern latitudes.

- Pointing:
 - Collaborative prime/rider coordination designs
 - ISS_159TI_HIGHRES001_PRIME observation w/ VIMS (-02:15 to +02:15)
 - No custom period
 - RBOT-friendly secondaries used in all cases except for RADAR and CIRS limb observations.
 - Secondaries are flexible except in the following cases:
 - VIMS_159TI_STELLOCC001_PRIME: Significant science impact if secondary changed (occultation)
- Data Volume:
 - No issues.
- DSN:
 - No special requests.
- Opmodes:
 - No special op modes.
- Hydrazine:
 - No thrusters.
- Special Activities:
 - None.

Sequence Liens:

None.

Segment Checklist p1

- TOST rev 159 T80

Item	Disposition notes, or X if complete
1. Disposition all requests in CIMS - approve all pending requests, no outstanding revisions/new requests	Х
2. No rocking downlinks. No AZSCANS (IGAPIMAGE). No arrayed downlinks.	Х
3. Examine SPASS, ensure SP turns correctly designated PRIME or NEW WAYPOINT. Prime RSS observations require the Xband to Earth attitude be a waypoint, use DLTURN with spass type New Waypoint (also for DLTURN before Ybiases)	Х
4. Waypoints and downlinks are violation free (per CTV). NOTE ON ISSUES PAGE if periods of no valid waypoint	Х
5. SP turns have been checked and are violation free- use ctv_batch or PDT. Fix any issues found. First turn of segment has been checked using correct final attitude of previous segment. All turns use the slower XM slew rates and include 2 minutes turn margin. Allow extra turn time whenever possible to aid possible RBOT changes.	Х
6. YBIAS windows have been included as required, guidelines met per https://cassini.jpl.nasa.gov/sp/xxmdev/ybias_mpforum.pdf	Х
7. There are no more than 3 waypoint changes in a 24 hour period (DLTURN waypoints for YBIAS do not count)	Х
8. The minimum prime instrument request duration outside ±5 hours from a targeted satellite flyby is 30 minutes	Х
9. Custom handoffs are limited to ±3 hours around a targeted Titan flyby or an asymmetric 10 hour window for Icy Satellite flybys. Custom periods 1) designated properly with SPASS notes 2) requests have "pick up at" and "hand off at" information filled in correctly 3) turn times and handoff attitudes have been verified – early PDT work recommended!	N/A
10. PIEs are properly identified via _PIE naming convention. All agreed to PIEs have been integrated.	N/A
11. Prime/rider coordination: secondaries have all been reviewed and agreed to, collaborative observations are so designated, predesigned in PDT, prime instrument agrees to work with riders for collaborate designs	Х
12. Use rolling_sru if required. Follow rolling guidelines per SCO, see the ScoRules wiki page (linked to integration procedure)	Х
13. The secondary axis for downlinks that contain prime and backup OTMs is the same, and inertially fixed	N/A
14. Downlinks that contain OTPs only roll for the first 4 hours of the downlink pass max. OTB: Full rolling OK, unless SRU issues, then 4_Hr_Rolling max (NO split rolls)	N/A
15. There is one downlink pass block per OTM prime or backup window (one wedding cake for a split pass). Exception - if first split downlink pass is ≤4 hours can use 2 cakes, put playback_gap in 2nd pass, put OTP/OTB in name of BOTH passes (for CDA). MUST have a full length 9 hour station requested for NAV tracking data	Х

Segment Checklist p2

TOST rev 159 T80 Disposition notes. Item or X if complete 16. Moving any downlink pass to a different view period requires coordination with Navigation. Changes to the DSN strawman plan N/A require SPST manager approval. 17. Multi-revolution turns about the X-axis have an offset greater than or equal to 30 degrees N/A 18. Live moveable blocks (LMBs) include the appropriate time margin specified as a DEADTIME request in CIMS at the beginning and N/A end of the moveable block. TLM modes in separate OBSMOV request (n/a for RSS). Waypoint same entering as leaving, and is valid throughout. Avoid skeet shoots in LMBs. If CMT management required, contain within LMB. Live moveable blocks use an LMB epoch and use the appropriate epoch naming conventions. Live Update Blocks use a LUB epoch (RSS only). 19. Pointing is not altered for science during any SCO/MP activity that has pointing requirements (e.g., dust hazards). [Note that science N/A turns are allowed for all but the first minute of an inbound thruster transition during a Titan or icy satellite flyby. No science turns are allowed during any portion of the outbound transition] 20. All stellar occultation observations include an additional +/-20 minutes of time (40 minutes total) when they occur within -1 day to +2 Χ days of Saturn periapse 21. All Ground and Live Moveable blocks associated with non-targeted geometric events (e.g., solar and earth occultations) include an Χ additional +/-20 minutes of time margin (40 minutes total) to account for reference trajectory changes. 22. Check your GMB, LMB, LUB, Occ times against current reference trajectory (Tour Atlas) Χ 23. Dual playback of high value data is performed within this segment and does not affect downstream segments. CIMS entries are N/A correct and SPASS type Note. SSR-A is emptied after the first downlink. Open a SPLAT item (tied to the ENGR request that resets the pointers, ie the DUALPB_CDS request) which says, "During DSN negotiations ensure that SSR-A is emptied before the pointers are reset. This item cannot be closed until the DSN negotiations are complete for both downlink passes, or the dual playback is deleted." Χ 24. Run the resource checker in CIMS and fix errors found. Remaining notes disposition here or on notes page 25. SMT: note if SSR not empty at end of segment, have approval from following segment. No carryover across sequence boundaries. Χ Aim for empty SSR every 4 days. No negative SSR margin during integration. List discrepancies on notes page. 26. Examine SMT warnings report, include dispositions here or on notes page of any items No data policing during part of RADAR warmup (this is ok) 27. RSS boresight: one _SP pass, two _PRIME downlink passes, one hour observation block in SNER_3 N/A

Segment Checklist p3

- TOST rev 159 T80

Item	Disposition notes, or X if complete
28. Examine "ap_downlink report check" output, include dispositions here or on notes page of any items (see next two items).	70M usage is 100%; # of seq passes <5 (both ok for TOST)
29. List any DSN stations requested during maintenance periods, AND JUSTIFICATION. AVOID!!!!!	Х
30. Avoid requesting two overlapping stations (except for RSS science) whenever possible – use RSS station for downlink too	Х
31. Compare RSS requests to DSN requests, make sure they jive (ORT, occ, etc), ORTs are integrated.	N/A
32. Apoapse segments only: List your percent 70M stations requested - avoid >35%.	N/A
33. Apoapse segments only: Follow Integration Guideline & Constaint #15c regarding "two out of three" types of science per RBOT segment. ME OTM's split an RBOT segment.	N/A
34. Periapse segments: >3 hr observations with >60 degree target motion are broken up by a 20 min inertial period (lien if not explicit in SPASS)	Х
35. Support images use _XXM or _XXM3 activity type	N/A
36. In CIMS check for "start before", "end before", "start after", "end after" requests - fix if any problems found	Х
37. Verify OPNAVs are in SNER5 and are support_image class, sanity check rest of tlm modes (RADAR 15 min in 5A/activity in 5A or 8, etc)	N/A
38. If sequence boundary at START of segment, ensure IVPGAP info correct, NO "start before" MAPS requests, OpNav is not first thing in segment	N/A
39. If sequence boundary at END of segment (ie in the next segment), ensure 6 "SEQ" upload DSN passes - will probably ripple into preceding segment(s), notify them. Last pass has Ybias window in front, no bonus science. NO "end after" MAPS requests	N/A
40. Verify opmodes correct (RSS and RADAR especially), teams going to sleep have agreed? MIMI: not in sleep during RPX? Use table at https://cassini.jpl.nasa.gov/wiki/bin/view/Cassini/XXMOpModes	Х
41. If conjunction is in your segment, see Conjunction page on SP Wiki	N/A

- TOST rev 159 T80

Item	Disposition notes, or X if complete
42. RAMAVOID: new waypoint, NOT in custom period	N/A
43. If on thrusters, confirm deadbands	N/A
44. Segment products linked to XXM deliveries page, & this package when you are done	Х